

## MENISCAL ALLOGRAFT TRANSPLANTATION REHABILITATION PROTOCOL

	<b>WEIGHT BEARING</b>	<b>BRACE</b>	<b>ROM</b>	<b>EXERCISES</b>
<b>PHASE I</b> 0-2 weeks	Heel touch weight bearing with crutches	Locked in full extension for sleeping and all activity  Off for exercises and hygiene	0-90° when non-weight bearing	Heel slides, quad sets, patellar mobs, SLR, SAQ**  <b>No weight bearing with flexion &gt;90°</b>
<b>PHASE II</b> 2-8 weeks	<b>2-6 weeks:</b> Heel touch WB with crutches  <b>6-8 weeks:</b> Progress to full WB	<b>2-8 weeks:</b> Locked 0-90°  Off at night  Discontinue brace at 8 weeks	As tolerated	Addition of heel raises, total gym (closed chain), terminal knee extensions**  Activities w/ brace until 6 weeks; then w/o brace as tolerated  <b>No weight bearing with flexion &gt;90°</b>
<b>PHASE III</b> 8-12 weeks	Full	None	Full	Progress closed chain activities  Begin hamstring work, lunges/leg press 0-90°, proprioception exercises, balance/core/hip/glutes  Begin stationary bike
<b>PHASE IV</b> 12-20 weeks	Full	None	Full	Progress Phase III exercises and functional activities: single leg balance, core, glutes, eccentric hamstrings, elliptical, and bike  Swimming okay at 16 wks
<b>PHASE V</b> > 20 weeks	Full	None	Full	Advance to sport-specific drills and running/jumping once cleared by MD

\*Brace may be removed for sleeping after first post-operative visit (day 7-10)

\*\*Avoid any tibial rotation for 8 weeks to protect meniscus